

# Look-up tables of ground-based UV-vis stratospheric NO<sub>2</sub> column averaging kernels

François Hendrick and Michel Van Roozendael (BIRA-IASB)

Contact: [franch@oma.be](mailto:franch@oma.be)

## 1. Introduction

A climatology of stratospheric NO<sub>2</sub> column averaging kernels has been developed with the aim to consolidate and better characterize the time-series of twilight zenith-sky NO<sub>2</sub> measurements produced by UV-vis and SAOZ spectrometers from the NDACC network. Column averaging kernels are needed for satellite nadir NO<sub>2</sub> validation in order to compare the sensitivity of the ground-based and space-borne instruments to the NO<sub>2</sub> vertical profile. Enhancing the usability of the correlative data sets is also the main reason why standard guidelines for data reporting like GEOMS (Global Earth Observation Metadata Standard; see <http://avdc.gsfc.nasa.gov/index.php?site=1178067684>) recommend to include metadata like averaging kernels in data files. We describe a new multi-entry database of ground-based UV-vis NO<sub>2</sub> column averaging kernels applicable at the global scale. It is complementary to the NO<sub>2</sub> air mass factor (AMF) climatology recently developed in order to homogenize the NDACC UV-vis stratospheric NO<sub>2</sub> column data sets.

## 2. Description

The proposed database of stratospheric NO<sub>2</sub> column averaging kernels is based on the theoretical work of *Eskes and Boersma (2003)* and has been generated using the UVSPEC/DISORT radiative transfer model (*Hendrick et al., 2006*) and the harmonic climatology of stratospheric NO<sub>2</sub> profile developed by *Lambert et al. (1999, 2000)*. This climatology consists of a Fourier harmonic decomposition of UARS HALOE v19 and SPOT-4 POAM-III v2 NO<sub>2</sub> profile data records. It covers the 20-60 km altitude range for the present application. Between 12 and 17 km, the NO<sub>2</sub> profiles are complemented by a climatology derived from SAOZ balloon observations (*F. Goutail, personal communication*). The NO<sub>2</sub> concentration is set to zero below 12 km altitude. In addition to the NO<sub>2</sub> profile climatology, the parameters considered in building the look-up tables (LUTs) are: wavelength, ground albedo, and SZA. Table 1 summarizes these different parameters and their corresponding values.

Parameter	Values
Lambert et al.'s NO <sub>2</sub> profile climatology	<ul style="list-style-type: none"><li>- Latitude: 85°S to 85°N step 10°</li><li>- Month: 1 (Jan) to 12 (Dec) step 1</li><li>- Sunrise and sunset</li><li>- Altitude range : 20-60 km</li></ul>
SAOZ balloon NO <sub>2</sub> profile climatology	<ul style="list-style-type: none"><li>- Latitude: tropics, mid- and high-latitudes</li><li>- Resolved in seasons (spring, summer, fall, winter)</li><li>- Altitude range: 12-20 km</li></ul>
Wavelength	350 to 550 nm step 40 nm

Surface albedo	0 and 1
SZA	90° (SZA representative of ground-based UV-vis measurements at twilight)
Aerosols	Aerosol extinction profile corresponding to background conditions and constructed from the aerosol model of <i>Shettle (1989)</i> included in UVSPEC/DISORT

Table 1: Parameters of the look-up table and their corresponding values.

The NO<sub>2</sub> column averaging kernels climatology consists of 2 look-up tables (one for sunrise and one for sunset conditions; size: 1 MB each). An interpolation routine has been developed in order to extract appropriately parameterized NO<sub>2</sub> column averaging kernels for the different NDACC stations. This routine is written in FORTRAN 77 and a DOS executable has been created. The source code is also available for compilation on LINUX machines. In addition, a global monthly climatology of the surface albedo at two wavelengths (380 and 494 nm) is coupled to the interpolation routine so that realistic albedo values can be obtained in a transparent way. This albedo climatology is extracted from the GOME surface albedo database developed by *Koelemeijer et al.* (2003). It consists of 24 (12x2) look-up tables, one for each month of the year and each wavelength. Albedo values are given for grid-cells of 1° x 1° (latitude: -89.5° to 89.5°; longitude: -179.5° to 179.5°).

### 3. How to use the NO<sub>2</sub> column averaging kernel climatology?

The zip file contains 31 files: 2 NO<sub>2</sub> column averaging kernel look-up tables, 24 surface albedo look-up tables (size: 1.1 MB each), a DOS executable ('no2\_avk\_interpolation\_v1\_0\_dos'), the source code in FORTRAN 77 ('no2\_avk\_interpolation\_v1\_0.for'), two input files for selecting parameter values, and one output file. All the files should be located in the same directory for a proper use of the climatology. In the file 'input\_file\_no2\_avk.dat', the user can enter values for wavelength, latitude, longitude, and surface albedo. Regarding the albedo, the user must give a value to a flag in order to determine whether he wants to use the albedo climatology (flag=1) or not (flag=2). The user has also to define the name of the file (here called 'DAY\_FILE.dat'; maximum number of lines in this file: 500000) containing lines with year, decimal day numbers (decimals being decimal UT time). The resulting NO<sub>2</sub> column averaging kernels are stored in the output file called 'no2\_avk\_output.dat'.

### 4. References

- Eskes, H. J., and K. F. Boersma, Averaging kernels for DOAS total-column satellite retrievals, *Atmos. Chem. Phys.*, 3, 1285-1291, 2003.
- Hendrick, F., M. Van Roozendaal, A. Kylling, A. Petritoli, S. Sanghavi, R. Schofield, C. von Friedeburg, F. Wittrock, and M. De Mazière, Intercomparison exercise between different radiative transfer models used for the interpretation of ground-based zenith-sky and multi-axis DOAS observations, *Atm. Chem. Phys.*, 6, 93-108, 2006.

- Koelemeijer, R. B. A., J. F. de Haan, and P. Stammes, A database of spectral surface reflectivity in the range 335-772 nm derived from 5.5 years of GOME observations, *J. Geophys. Res.*, 108 (D2), doi 10.1029/2002JD002429, 2003
- Lambert, J.-C., J. Granville, M. Van Roozendael, A. Sarkissian, F. Goutail, J.-F. Müller, J.-P. Pommereau, and J.M. Russell III, A climatology of NO<sub>2</sub> profile for improved Air Mass Factors for ground-based vertical column measurements, in *Stratospheric Ozone 1999*, N.R.P. Harris, M. Guirlet, and G.T. Amanatidis (Eds.), Air Pollution Research Report 73 (CEC DG XII), pp. 703-706, 1999.
- Lambert, J.-C., J. Granville, M. Van Roozendael, J.-F. Müller, F. Goutail, J.-P. Pommereau, A. Sarkissian, P. V. Johnston, and J. M. Russell III, Global Behaviour of Atmospheric NO<sub>2</sub> as Derived from the Integrated Use of Satellite, Ground-based Network and Balloon Observations, in *Atmospheric Ozone - 19th Quad. Ozone Symp.*, Sapporo, Japan, 2000, Ed. by NASDA, pp. 201-202, 2000.
- Shettle, E. P., Models of aerosols, clouds, and precipitation for atmospheric propagation studies, AGARD Conference Proceedings No. 454: Atmospheric propagation in the UV, visible, IR and mm-region and related system aspects, 1989.

## 5. Contact

For questions, comments or bug report regarding the stratospheric NO<sub>2</sub> averaging kernels climatology, please contact François Hendrick at the Belgian Institute for Space Aeronomy (IASB-BIRA).

E-mail: franch@oma.be